



TETRIS EW

BEFORE USING THIS PRODUCT, READ THE LABEL CAREFULLY

TETRIS EW

FUNGICIDE

Reg. No. L8773, Act/Wet No. 36 of/van 1947 1:07/09/2011

An oil-in-water emulsion, systemic fungicide for the control of powdery mildew on crops as listed.

'n Olie-in-water emulsie, sistemiese swamdoder vir die beheer van poeieragtige skimmel op gewasse soos aangedui.

FRAC FUNGICIDE GROUP CODE 3 FRAC SWAMDODERGROEKODE

ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL

Tetraconazole (triazole) 100 g/l Tetrakonasool (triasool)

Registration Holder / Registrasiehouer
Meridian Agrochemical Company (Pty) Ltd. / (Edms.) Bpk.
Reg. No. 2000/010819/07
P.O. Box / Posbus 436, Modderfontein, 1645, South Africa / Suid-Afrika

5l

UN Number: 3082

DIRECTIONS FOR USE ENCLOSED
Batch Number
Date of manufacture
REFER TO DETAILS AS PRINTED
ON CONTAINER / BAG

Meridian agritech

CAUTION / VERSIGTIG

WARNINGS:

The following number of days should be allowed between last application and harvest of the crop:

Wine and Table grapes 14 days

- Handle with care.
- May irritate eyes and cause moderate skin irritation.
- Store in a cool, dry place in original, tightly closed containers in a well ventilated area.
- Avoid temperatures below 0 °C and above 35 °C.
- Store away from food and feed.
- Keep out of reach of children, uninformed persons and animals.
- FLAMMABLE - Keep away from fire and sparks.
- Do not use on hydroponically or container grown crops.
- Toxic to birds, fish and other aquatic organisms.
- **TETRIS EW** has been tested on some of the most important commercial cultivars without any phytotoxic effects. More susceptible cultivars may be released in the future which need to be tested; for this apply **TETRIS EW** to a limited test area before commercial applications are made.

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions; quality of dilution water; compatibility with other substances not indicated on the label and the occurrence of resistance of the disease organisms against the remedy concerned as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS:

- Avoid inhalation of spray mist and fumes.
- Avoid contact with skin and eyes.
- In case of skin contact, wash thoroughly with soap and water.
- Wash contaminated clothing.
- Do not eat, drink or smoke whilst applying or preparing the spray mix, or before washing face and hands.
- Prevent spray drift onto other crops, grazing, rivers, dams, boreholes and any area not under treatment.
- Prevent contamination of food, feed, drinking water and eating utensils.
- Clean applicator after use. Dispose of wash water where it will not contaminate crops, grazing, rivers, dams etc.
- Rinse the empty container three times with a volume of clean water equal to a minimum of 10 % of the container. Add the rinsate to the content of the spray tank before destroying the container.
- Destroy empty container and do not use it for any other purpose.
- Do not apply **TETRIS EW** to any crop or situation not mentioned in this label.
- Do not apply this product in combination or tank mixture with any other product or agricultural remedy which is not specifically listed on this label. Meridian Agrochemical Company (Pty) Ltd accepts no responsibility for any negative effects experienced if the instructions on this label are not adhered to, unless a Meridian representative was consulted beforehand.
- If **TETRIS EW** is used in a tank mix with other recommended products, consideration should be given to the restrictions of those respective products.

RESISTANCE WARNING:

TETRIS EW contains tetriconazole (triazole), a fungicide with specific activity against powdery mildew on grapes. For resistance management, **TETRIS EW** is a FRAC group code 3 fungicide. Any fungus population may contain individuals naturally resistant to **TETRIS EW** and other group code 3 fungicides. The resistant individuals can eventually dominate the disease population if these fungicides are used repeatedly. These resistant fungi may not be controlled by **TETRIS EW** or any other group code 3 fungicide. To prevent pathogen populations from becoming resistant to **TETRIS EW** it is essential that the following guidelines be followed:

- Avoid exclusive, repeated use of fungicides from the same fungicide group code.
- Alternate **TETRIS EW** with other registered fungicides with a different mode of action.
- Do not make more than four **TETRIS EW** applications to the same crop in one season.
- Use the recommended rate on the label. Do not reduce rates.
- Do not exceed recommended spray intervals.

Since the occurrence of resistance cannot be forecast, users are advised to keep treated crops under close observation. If treatment is not effective following the use of **TETRIS EC** as recommended, a resistant strain may be present. If a resistant strain is positively identified, consideration should be given to prompt use of a fungicide with a different mode of action for which there is no record of resistance to the relevant disease. Meridian Agrochemical Company (Pty) Ltd will not accept any liability for performance failures in the event of a build-up of disease resistance resulting from inadequate resistance management practices as indicated on the label. For specific information on resistance management, contact the registration holder of this product or visit the FRAC website at <http://www.frac.info/frac/index.html>.

DIRECTIONS FOR USE

Use only as directed.

General Properties:

TETRIS EW is a systemic fungicide with excellent protective properties against powdery mildew (*Oidium tuckeri*) on table and wine grapes.

Compatibility:

The compatibility of **TETRIS EW** with other products may be influenced by the formulation of the product involved as well as the quality of the water. Since the formulation of other products may change and water quality may vary from farm to farm, a physical compatibility test should always be carried out prior to application.

Mixing Instructions:

Half fill the spray tank with clean water and whilst agitating, pour the required amount of **TETRIS EW** into the spray tank and then top up the tank to the required water volume. Ensure thorough agitation during filling and spraying.

Application:

TETRIS EW may be applied with conventional high volume spray equipment. Calibrate the apparatus before application to ensure that the correct dosage is applied.

CROP AND DISEASE	APPLICATION VOLUME	REMARKS
TABLE AND WINE GRAPES Powdery mildew (<i>Oidium tuckeri</i>)	30 mL / 100 ℥ water every 10 – 14 days OR 45 mL / 100 ℥ water every 18 – 21 days	TETRIS EW is most effective when applied preventively or at the early stages of infection, before symptoms are visible on the plants. Commence spraying when the first shoots are 100 – 250 mm long and repeat the treatment at intervals as indicated in the 'Remarks' below. Do not exceed six applications

		See additional 'Remarks' below.
--	--	---------------------------------

Remarks:

- Always use the higher dosage rate if spray interval exceeds 14 days. Under conditions favouring disease development, spray intervals should not exceed 14 days.
- Gradually increase the spray volume from 250 l to 1200 l per ha depending on the vineyard size and density. Ensure thorough coverage of leaves and bunches.
- Apply **TETRIS EW** in a programme spray schedule as a full cover spray to the point of run-off.

WAARSKUWINGS:

Die volgende aantal dae moet toegelaat word tussen die laaste toediening en oes van die gewas:

Wyn- en Tafeldruwe 14 dae

- Hanteer met sorg.
- Mag oë irriteer en matige velirritasie veroorsaak.
- Bê in 'n koel, droë plek in die oorspronklike dig toegemaakte houers in 'n goed geventileerde area.
- Vermy temperature onder 0 °C en bokant 35 °C.
- Bêre weg van voedsel en voer.
- Hou buite bereik van kinders, oningesigte persone en diere.
- VLAMBAAR - Hou weg van vuur en vonke.
- Moet nie op hidroponeise of potgekweekte gewasse gebruik nie.
- Giftig vir voëls, visse en ander waterorganismes.
- **TETRIS EW** is sonder enige fitotoksisiteitskade op die mees belangrikste kommersiële kultivars getoets. Meer vatbare kultivars mag in die toekoms vrygestel word wat eers getoets moet word; hiervoor moet **TETRIS EW** eers op 'n klein toetsoppervlakte aangewend word, alvorens groot kommersiële bespuitings gemaak word.

Alhoewel hierdie middel omvattend onder 'n groot verskeidenheid toestande getoets is, waarborg die registrasiehouer nie dat dit onder alle toestande doeltreffend sal wees nie aangesien die werking en effek daarvan beïnvloed kan word deur faktore soos abnormale klimaats-en bergingstoestande, kwaliteit van verdunningswater verenigbaarheid met ander stowwe wat nie op die etiket aangedui is nie en die voorkoms van weerstand van die siekte organismes teen die betrokke middel sowel as die metode, tyd en akkuraatheid van toediening. Die registrasiehouer aanvaar geen verantwoordelikheid vir skade aan gewasse, plantegroei, die omgewing of vir nadelige effek op mens of dier vir 'n gebrek aan prestatie van die betrokke middel betrokke as gevolg van versuim van die gebruiker om die etiketaanwysings na te volg of die ontstaan van toestande wat nie in terme van die registrasie voorsien is. Raadpleeg die verskaffer in die geval van enige onsekerheid.

VOORSORGMAATREËLS:

- Vermy inaseming van spuitmis en dampe.
- Vermy kontak met vel en oë.
- In die geval van kontak met die vel was deeglik met seep en water.
- Was besoedelde klere.
- Moet nie eet, drink of rook tydens toediening of die voorbereiding van die spuitmengsel, of voordat jou gesig en hande gewas is nie.
- Voorkom oorwaai van spuitnewel na ander gewasse, weiding, riviere, damme, boorgate en enige gebied nie onder behandeling nie.
- Voorkom besoedeling van voedsel, voer, drinkwater en eetgerei.
- Maak toedieningsapparaat deeglik skoon na gebruik. Raak ontslae van waswater waar dit nie gewasse, weiding, riviere, damme, ens. kan besoedel nie.
- Spoel die leë houer **drie keer** uit met 'n volume skoon water gelykstaande aan 'n minimum van 10% van die houer. Gooi die spoelwater by die inhoud van die spuittenk voordat die houer vernietig word.
- Vernietig leë houer en moenie dit vir enige ander doel gebruik nie.
- Moet nie **TETRIS EW** toedien op enige gewas of situasie wat nie op hierdie etiket verskyn nie.
- Moet nie hierdie produk in kombinasie of tenkmengsel saam met enige ander produk of landboumiddel gebruik, wat nie spesifiek op hierdie etiket gelys word nie. Meridian landbouchemiese Maatskappy (Edms.) Bpk. aanvaar geen verantwoordelikheid vir enige negatiewe gevolge ervaar indien die instruksies op die etiket nie nagekom word nie, tensy 'n Meridian verteenwoordiger vooraf geraadpleeg is.
- As **TETRIS EW** in 'n tenkmengsel met ander aanbevole produkte gebruik word, moet

oorweging geskenk word aan die beperkings van daardie betrokke produkte.

WEERSTANDSWAARSKUWING:

Tetris EW bevat Tetrakonasool (Triasool), 'n swamdoder met spesifieke werking teen poeieragtige skimmel op druwe. Vir weerstandsbestuur is **TETRIS EW** 'n FRAC groepkode 3 swamdoder. Enige swam mag individue bevat met 'n natuurlike weerstand teen **TETRIS EW** of enige ander groepkode 3 swamdoder het. Die weerstandbiedende individue kan uiteindelik die siekte-bevolking oorheers as hierdie swamdoders herhaaldelik gebruik word. Hierdie weerstandbiedende swamme sal waarskynlik nie deur **TETRIS EW** of enige ander groepkode 3 swamdoder beheer word nie. Om patogeen-populasies te verhoed om teen **TETRIS EW** weerstandbiedend te raak, is dit noodsaaklik dat die volgende riglyne gevolg word:

- Vermy die eksklusieve herhaaldelike gebruik van swamdoders met dieselfde groepkode.
- Wissel **TETRIS EW** af met ander geregistreerde swamdoders met 'n ander metode van werking.
- Moenie meer as vier **TETRIS EW** toedienings op dieselfde gewas in een seisoen maak nie.
- Gebruik die aanbevole dosis op die etiket. Moenie dosisse verminder nie.
- Moenie aanbevole sputintervalle oorskry nie.

Aangesien die voorkoms van weerstand nie voorspel kan word nie, word gebruikers aangeraai om behandelde gewasse noukeurig dop te hou. Indien die behandeling met **TETRIS EW** soos aanbeveel nie doeltreffend was nie, mag 'n weerstandbiedende-populasie teenwoordig wees. As 'n weerstandbiedende-populasie positief geïdentifiseer word, moet daar oorweging geskenk word aan die gebruik van 'n swamdoder met 'n ander metode van werking waarvoor daar geen rekord van weerstand teen die betrokke siekte is nie. Meridian Agrochemical Company (Edms.) Bpk. sal geen aanspreeklikheid vir enige gebrek aan prestasie in die geval van 'n opbou van weerstand teen siektes as gevolg van onvoldoende weerstandsbesturspraktyke aanvaar soos aangedui op die etiket. Vir spesifieke inligting oor weerstandsbestuur kontak die registrasiehouer van hierdie produk of besoek die FRAC webwerf by <http://www.frac.info/frac/index.html>.

GEBRUIKSAANWYSINGS

Gebruik slegs soos aangedui.

Algemene eienskappe:

Tetris EW is 'n sistemiese swamdoder met uitstekende beskermende eienskappe teen poeieragtige skimmel (*Oidium tuckeri*) op tafeldruwe. **Tetris EW** is die mees doeltreffenste wanneer dit voorkomend toegedien word, of by die vroeë stadiums van die infeksie, voor simptome op die plante sigbaar is.

VERENIGBAARHEID

Die verenigbaarheid van **Tetris EW** met ander produkte kan beïnvloed word deur die formulering van die betrokke produk, sowel as die kwaliteit van die sputtwater. Aangesien die formulering van ander produkte kan verander, en die kwaliteit van die water kan verander van plaas tot plaas, moet 'n fisiese verenigbaarheidstoets altyd uitgevoer word, voordat gespuit word.

MENGINSTRUKSIES

Maak die sputtenk halfvol met skoon water terwyl dit geroer word, voeg die benodigde hoeveelheid **Tetris EW** in die sputtenk en vul op met die verlangde volume water. Verseker deeglike roering tydens volmaak en toediening.

TOEDIENING

Tetris EW kan met konvensionele hoë-volume sputtoerusting toegedien word. Kalibreer die

apparaat voor toediening om te verseker dat die korrekte dosis toegedien word.

AANBEVELINGS

GEWAS EN SIEKTE	DOSIS/ha of soos aangedui	OPMERKINGS
TAFEL- & WYNDRUIWE Poeieragtige skimmel (<i>Oidium tuckeri</i>)	30 mL / 100 L water elke 10 – 14 dae OF 45 mL / 100 L water elke 18 – 21 dae	Begin sput wanneer die eerste lote 100-250 mm lank is en herhaal behandeling met tussenposes soos aangedui in die 'Notas' hieronder. Moet nie meer as ses toedienings maak nie. Sien bykomende 'Notas' hieronder.

Notas:

- Gebruik altyd die hoër dosis as sputintervalle meer as 14 dae oorskry. Onder toestande wat gunstig is vir die ontwikkeling van die siekte, moet die sputintervalle nie langer as 14 dae wees nie.
- Verhoog geleidelik die sputvolume van 250 L tot 1200 L per ha, afhangende van die wingerd se grootte en digtheid. Verseker deeglike bedekking van blare en trosse.
- Dien **Tetris EW** toe in 'n programbespuiting-skedule as 'n volle dekbespuiting tot by die punt van afloop.